

B5

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number
WO 01/73029 A2(51) International Patent Classification⁷: C12N 15/12,
C07K 14/705, 14/715, 16/28, A01K 67/027, C12N 15/63,
5/10, C12P 21/00, G01N 33/53, C12Q 1/68, G01N 33/50

(21) International Application Number: PCT/US01/09522

(22) International Filing Date: 27 March 2001 (27.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/192,419 27 March 2000 (27.03.2000) US
60/230,459 6 September 2000 (06.09.2000) US
09/666,535 20 September 2000 (20.09.2000) US(71) Applicant: PE CORPORATION (NY) [US/US]; Robert
A. Millman, 761 Main Avenue, Norwalk, CT 06859 (US).(72) Inventors: YE, Jane; c/o Celera, 45 West Gude Drive,
Rockville, MD 20850 (US). CRAYCHIK, Anibal; c/o
Celera, 45 West Gude Drive, Rockville, MD 20850 (US).
DI FRANCESCO, Valentina; c/o Celera, 45 West Gude
Drive, Rockville, MD 20850 (US). BEASLEY, Ellen, M.;
c/o Celera, 45 West Gude Drive, Rockville, MD 20850
(US).(74) Agent: MILLMAN, Robert, A.; Celera Genomics Corp.,
45 West Gude Drive C2-4, Rockville, MD 20850 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

[Continued on next page]

(54) Title: ISOLATED HUMAN G-PROTEIN COUPLED RECEPTORS, NUCLEIC ACID MOLECULES ENCODING HUMAN
GPCR PROTEINS, AND USES THEREOF

```

1 ATGTACAACG GGTGCTGCTG CGGCATCGAG GGGGACACCA TCTCCAGGT
51 GATGCCGCCG CTGCTCATTT TGGCCCTTGT CTTGGGCGCA CTAGGCAATG
101 GGGTGGCCCT GTGTGGTTTC TGCTTCCACA TGAAGACCTG GAAGCCGAGC
151 ACTGTTTACC TTTTCAATTT GGGCGTGGCT GATTTCCTCC TTATGATCTG
201 CTTGCTTTT CGGACAGACT ATTACCTCAG ACGTAGACAC TGGGCTTTTG
251 GGGACATTCG CTGGCGAGTG GGGCTCTTCA CTTGGGCGAT GAAGAGCGCC
301 GGGAGCATCC TGTCTCTTAC GCTGCTGCTT GCGGACAGCT ATTTCAAGGT
351 GGTCCACCCC CACCCGCGGG TGAACACTAT TCCACCCCGG GTGGCGGCTG
401 GCATGCTCTG CACCCGTGTTG GGCCTGGTCA TCTTGGGAGC AGTGTATCTT
451 TTGCTGGAGA ACCATCTCTG CGTGCAAGAG ACGGCCGTCT CTTGTGAGAG
501 CTTGATCATG GAGTGGGCGA ATGGCTGSCA TGACATCATG TTCCAGCTGG
551 AGTTCTTTAT GCGGCTGGGC ATCATCTTAT TTGCTCCTTT CAAGATTGTT
601 TGGAGCCTGA GCGCGAGGCA GCGAGTGGCC AGACAGGCTC GGTGAGAA
651 GGGGACCCGG TTCATCATGG TGGTGGCAAT TGTGTTCATC ACATGCTAGC
701 TGGCAGCGGT GTCTGCTAGA CTCTATTTCG TCTGACCGGT GCGCTGAGT
751 GCGTGGGATC CCTGTGTCGA TGGGGCCCTG CACCTAAGCC TGAAGCTTCA
801 CTACATGAAC AGCATGCTGG ATCCCTGGGT GTATTATTTT TCAAGCCCTT
851 CTTTTCACAA ATTCTAGAAC AAGCTCAAAA TCTGGAGTCT GAAAGCCGAG
901 GAGCCAGGAC ACTCAAAAC ACAAAGGCGG GAAGAGATGC CAATTTGCAA
951 CCTCGGTGCG AGGACTTGCA TCAGTGTGGC AAATAGTTTC CAAAGCCAGT
1001 CTGATGGGCA ATGGGATCCC CACATTGTTG AGTGGCACTG A

```

FEATURES:
Start: 1
Stop: 1039

HOMOLOGOUS PROTEIN:
Top 10 BLAST Hits:

gi15174461 ref NP_006009.1 putative chemokine receptor; GTP-b...	344	7e-94
gi14595317 ref NP_005290.1 G protein-coupled receptor 31 >gil...	175	6e-43
gi16716509 gb AA22668.1 AF140708.1 (AF140708): G protein couple...	172	4e-42
gi1464327 sp P34956 P2YR_CHICK P2Y PURINOCEPTOR 1 (ATP RECEPTOR...	138	6e-32
gi11352693 sp P49652 P2YR_MELGA P2Y PURINOCEPTOR 1 (ATP RECEPTO...	138	6e-32
gi11352691 sp P48042 P2YR_BOVIN P2Y PURINOCEPTOR 1 (ATP RECEPTO...	138	1e-31
gi14505557 ref NP_002554.1 purinergic receptor P2Y, G-protein...	135	5e-31
gi11352695 sp P49651 P2YR_RAT P2Y PURINOCEPTOR 1 (ATP RECEPTOR)...	135	5e-31
gi16075193 ref NP_032756.1 purinergic receptor P2Y, G-protein...	131	4e-30
gi14885311 ref NP_005287.1 G protein-coupled receptor 22 >gil...	120	2e-29

EXPRESSION INFORMATION FOR MODULATORY USE (from cDNA panel screening):
Human Brain
Human Heart
Human Lung
Human Uterus
Human Placenta
Human Thyroid

(57) Abstract: The present invention provides amino acid sequences of peptides that are encoded by genes within the Human genome, the GPCR peptides of the present invention. The present invention specifically provides isolated peptide and nucleic acid molecules, methods of identifying orthologs and paralogs of the GPCR peptides and methods of identifying modulators of the GPCR peptides.

WO 01/73029 A2

28

WO 01/73029 A2



(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

-- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(aminergic GPCRs) represent a large group of GPCRs that share a common evolutionary ancestor and which are present in both vertebrate (deuterostome), and invertebrate (protostome) lineages. This family of GPCRs includes, but is not limited to the 5-HT-like, the dopamine-like, the acetylcholine-like, the adrenaline-like and the melatonin-like GPCRs.

5 Dopamine receptors

The understanding of the dopaminergic system relevance in brain function and disease developed several decades ago from three diverse observations following drug treatments. These were the observations that dopamine replacement therapy improved Parkinson's disease symptoms, depletion of dopamine and other catecholamines by reserpine caused depression and antipsychotic
10 drugs blocked dopamine receptors. The finding that the dopamine receptor binding affinities of typical antipsychotic drugs correlate with their clinical potency led to the dopamine overactivity hypothesis of schizophrenia (Snyder, S.H., *Am J Psychiatry* 133, 197-202 (1976); Seeman, P. and Lee, T., *Science* 188, 1217-9 (1975)). Today, dopamine receptors are crucial targets in the pharmacological therapy of schizophrenia, Parkinson's disease, Tourette's syndrome, tardive dyskinesia and Huntington's disease.
15 The dopaminergic system includes the nigrostriatal, mesocorticolimbic and tuberoinfundibular pathways. The nigrostriatal pathway is part of the striatal motor system and its degeneration leads to Parkinson's disease; the mesocorticolimbic pathway plays a key role in reinforcement and in emotional expression and is the desired site of action of antipsychotic drugs; the tuberoinfundibular pathways regulates prolactin secretion from the pituitary.

20 Dopamine receptors are members of the G protein coupled receptor superfamily, a large group proteins that share a seven helical membrane-spanning structure and transduce signals through coupling to heterotrimeric guanine nucleotide-binding regulatory proteins (G proteins). Dopamine receptors are classified into subfamilies: D1-like (D1 and D5) and D2-like (D2, D3 and D4) based on their different ligand binding profiles, signal transduction properties, sequence homologies and
25 genomic organizations (Civelli, O., Bunzow, J.R. and Grandy, D.K., *Annu Rev Pharmacol Toxicol* 33, 281-307 (1993)). The D1-like receptors, D1 and D5, stimulate cAMP synthesis through coupling with Gs-like proteins and their genes do not contain introns within their protein coding regions. On the other hand, the D2-like receptors, D2, D3 and D4, inhibit cAMP synthesis through their interaction with Gi-like proteins and share a similar genomic organization which includes introns within their
30 protein coding regions.

Serotonin receptors

Serotonin (5-Hydroxytryptamine; 5-HT) was first isolated from blood serum, where it was shown to promote vasoconstriction (Rapport, M.M., Green, A.A. and Page, I.H., *J Biol Chem* 176,

1243-1251 (1948). Interest on a possible relationship between 5-HT and psychiatric disease was spurred by the observations that hallucinogens such as LSD and psilocybin inhibit the actions of 5-HT on smooth muscle preparations (Gaddum, J.H. and Hameed, K.A., *Br J Pharmacol* 9, 240-248 (1954)). This observation lead to the hypothesis that brain 5-HT activity might be altered in psychiatric disorders (Wooley, D.W. and Shaw, E., *Proc Natl Acad Sci U S A* 40, 228-231 (1954); Gaddum, J.H. and Picarelli, Z.P., *Br J Pharmacol* 12, 323-328 (1957)). This hypothesis was strengthened by the introduction of tricyclic antidepressants and monoamine oxidase inhibitors for the treatment of major depression and the observation that those drugs affected noradrenaline and 5-HT metabolism. Today, drugs acting on the serotonergic system have been proved to be effective in the pharmacotherapy of psychiatric diseases such as depression, schizophrenia, obsessive-compulsive disorder, panic disorder, generalized anxiety disorder and social phobia as well as migraine, vomiting induced by cancer chemotherapy and gastric motility disorders.

Serotonin receptors represent a very large and diverse family of neurotransmitter receptors. To date thirteen 5-HT receptor proteins coupled to G proteins plus one ligand-gated ion channel receptor (5-HT₃) have been described in mammals. This receptor diversity is thought to reflect serotonin's ancient origin as a neurotransmitter and a hormone as well as the many different roles of 5-HT in mammals. The 5-HT receptors have been classified into seven subfamilies or groups according to their different ligand-binding affinity profiles, molecular structure and intracellular transduction mechanisms (Hoyer, D. et al., *Pharmacol. Rev.* 46, 157-203 (1994)).

20 Adrenergic GPCRs

The adrenergic receptors comprise one of the largest and most extensively characterized families within the G-protein coupled receptor "superfamily". This superfamily includes not only adrenergic receptors, but also muscarinic, cholinergic, dopaminergic, serotonergic, and histaminergic receptors. Numerous peptide receptors include glucagon, somatostatin, and vasopressin receptors, as well as sensory receptors for vision (rhodopsin), taste, and olfaction, also belong to this growing family. Despite the diversity of signalling molecules, G-protein coupled receptors all possess a similar overall primary structure, characterized by 7 putative membrane-spanning .alpha. helices (Probst et al., 1992). In the most basic sense, the adrenergic receptors are the physiological sites of action of the catecholamines, epinephrine and norepinephrine. Adrenergic receptors were initially classified as either .alpha. or .beta. by Ahlquist, who demonstrated that the order of potency for a series of agonists to evoke a physiological response was distinctly different at the 2 receptor subtypes (Ahlquist, 1948). Functionally, .alpha. adrenergic receptors were shown to control vasoconstriction, pupil dilation and uterine inhibition, while .beta. adrenergic receptors were implicated in vasorelaxation, myocardial stimulation and bronchodilation (Regan et al., 1990).

analyses suggest that the ancestor of these proteins originally developed in concert with complex body plans and nervous systems.

The GPCR protein superfamily can be divided into five families: Family I, receptors typified by rhodopsin and the β 2-purinergic receptor and currently represented by over 200 unique members (Dohlman *et al.*, *Annu. Rev. Biochem.* 60:653-688 (1991)); Family II, the parathyroid hormone/calcitonin/secretin receptor family (Juppner *et al.*, *Science* 254:1024-1026 (1991); Lin *et al.*, *Science* 254:1022-1024 (1991)); Family III, the metabotropic glutamate receptor family (Nakanishi, *Science* 258 597:603 (1992)); Family IV, the cAMP receptor family, important in the chemotaxis and development of *D. discoideum* (Klein *et al.*, *Science* 241:1467-1472 (1988)); and Family V, the fungal mating pheromone receptors such as STE2 (Kurjan, *Annu. Rev. Biochem.* 61:1097-1129 (1992)).

There are also a small number of other proteins that present seven putative hydrophobic segments and appear to be unrelated to GPCRs; they have not been shown to couple to G-proteins. *Drosophila* expresses a photoreceptor-specific protein, bride of sevenless (boss), a seven-transmembrane-segment protein that has been extensively studied and does not show evidence of being a GPCR (Hart *et al.*, *Proc. Natl. Acad. Sci. USA* 90:5047-5051 (1993)). The gene *frizzled* (*fz*) in *Drosophila* is also thought to be a protein with seven transmembrane segments. Like boss, *fz* has not been shown to couple to G-proteins (Vinson *et al.*, *Nature* 338:263-264 (1989)).

G proteins represent a family of heterotrimeric proteins composed of α , β and γ subunits, that bind guanine nucleotides. These proteins are usually linked to cell surface receptors, e.g., receptors containing seven transmembrane segments. Following ligand binding to the GPCR, a conformational change is transmitted to the G protein, which causes the α -subunit to exchange a bound GDP molecule for a GTP molecule and to dissociate from the $\beta\gamma$ -subunits. The GTP-bound form of the α -subunit typically functions as an effector-modulating moiety, leading to the production of second messengers, such as cAMP (e.g., by activation of adenylyl cyclase), diacylglycerol or inositol phosphates. Greater than 20 different types of α -subunits are known in humans. These subunits associate with a smaller pool of β and γ subunits. Examples of mammalian G proteins include Gi, Go, Gq, Gs and Gt. G proteins are described extensively in Lodish *et al.*, *Molecular Cell Biology*, (Scientific American Books Inc., New York, N.Y., 1995), the contents of which are incorporated herein by reference. GPCRs, G proteins and G protein-linked effector and second messenger systems have been reviewed in *The G-Protein Linked Receptor Fact Book*, Watson *et al.*, eds., Academic Press (1994).

Aminergic GPCRs

One family of the GPCRS, Family II, contains receptors for acetylcholine, catecholamine, and indoleamine ligands (hereafter referred to as biogenic amines). The biogenic amine receptors

ISOLATED HUMAN G-PROTEIN COUPLED RECEPTORS, NUCLEIC ACID MOLECULES ENCODING HUMAN GPCR PROTEINS, AND USES THEREOF

RELATED APPLICATIONS

5 The present application claims priority to U.S. Serial Nos. 60/192,419, filed March 27, 2000 (Atty. Docket CL000397-PROV), 60/230,459, filed September 6, 2000 (Atty. Docket CL000782-PROV), and 09/666,535, filed September 20, 2000 (Atty. Docket CL000782).

FIELD OF THE INVENTION

10 The present invention is in the field of G-Protein coupled receptors (GPCRs) that are related to the chemokine receptor subfamily, recombinant DNA molecules, and protein production. The present invention specifically provides novel GPCR peptides and proteins and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

15

BACKGROUND OF THE INVENTION

G-protein coupled receptors

G-protein coupled receptors (GPCRs) constitute a major class of proteins responsible for transducing a signal within a cell. GPCRs have three structural domains: an amino terminal
20 extracellular domain, a transmembrane domain containing seven transmembrane segments, three extracellular loops, and three intracellular loops, and a carboxy terminal intracellular domain. Upon binding of a ligand to an extracellular portion of a GPCR, a signal is transduced within the cell that results in a change in a biological or physiological property of the cell. GPCRs, along with G-proteins and effectors (intracellular enzymes and channels modulated by G-proteins), are the components of a
25 modular signaling system that connects the state of intracellular second messengers to extracellular inputs.

GPCR genes and gene-products are potential causative agents of disease (Spiegel *et al.*, *J. Clin. Invest.* 92:1119-1125 (1993); McKusick *et al.*, *J. Med. Genet.* 30:1-26 (1993)). Specific defects in the rhodopsin gene and the V2 vasopressin receptor gene have been shown to cause various forms of
30 retinitis pigmentosum (Nathans *et al.*, *Annu. Rev. Genet.* 26:403-424(1992)), and nephrogenic diabetes insipidus (Holtzman *et al.*, *Hum. Mol. Genet.* 2:1201-1204 (1993)). These receptors are of critical importance to both the central nervous system and peripheral physiological processes. Evolutionary

Eventually, pharmacologists realized that these responses resulted from activation of several distinct adrenergic receptor subtypes. β adrenergic receptors in the heart were defined as β .sub.1, while those in the lung and vasculature were termed β .sub.2 (Lands et al., 1967).

α Adrenergic receptors, meanwhile, were first classified based on their anatomical
5 location, as either pre or post-synaptic (α .sub.2 and α .sub.1, respectively) (Langer et al., 1974). This classification scheme was confounded, however, by the presence of α .sub.2 receptors in distinctly non-synaptic locations, such as platelets (Berthelsen and Pettinger, 1977). With the development of radioligand binding techniques, α adrenergic receptors could be distinguished pharmacologically based on their affinities for the antagonists prazosin or yohimbine
10 (Stark, 1981). Definitive evidence for adrenergic receptor subtypes, however, awaited purification and molecular cloning of adrenergic receptor subtypes. In 1986, the genes for the hamster β .sub.2 (Dickson et al., 1986) and turkey β .sub.1 adrenergic receptors (Yarden et al., 1986) were cloned and sequenced. Hydropathy analysis revealed that these proteins contain 7 hydrophobic domains similar to rhodopsin, the receptor for light. Since that time the adrenergic receptor family
15 has expanded to include 3 subtypes of β receptors (Emorine et al., 1989), 3 subtypes of α .sub.1 receptors (Schwinn et al., 1990), and 3 distinct types of β .sub.2 receptors (Lomasney et al., 1990).

The cloning, sequencing and expression of α receptor subtypes from animal tissues has led to the subclassification of the α 1 receptors into α 1d (formerly known as α 1a or
20 1a/1d), α 1b and α 1c (formerly known as α 1c) subtypes. Each α 1 receptor subtype exhibits its own pharmacologic and tissue specificities. The designation " α 1a" is the appellation recently approved by the IUPHAR Nomenclature Committee for the previously designated " α 1c" cloned subtype as outlined in the 1995 Receptor and Ion Channel Nomenclature Supplement (Watson and Girdlestone, 1995). The designation α 1a is used
25 throughout this application to refer to this subtype. At the same time, the receptor formerly designated α 1a was renamed α 1d. The new nomenclature is used throughout this application. Stable cell lines expressing these α 1 receptor subtypes are referred to herein; however, these cell lines were deposited with the American Type Culture Collection (ATCC) under the old nomenclature. For a review of the classification of α 1 adrenoceptor subtypes, see,
30 Martin C. Michel, et al., Naunyn-Schmiedeberg's Arch. Pharmacol. (1995) 352:1-10.

The differences in the α adrenergic receptor subtypes have relevance in pathophysiologic conditions. Benign prostatic hyperplasia, also known as benign prostatic hypertrophy or BPH, is an illness typically affecting men over fifty years of age, increasing in severity with increasing age. The symptoms of the condition include, but are not limited to,

increased difficulty in urination and sexual dysfunction. These symptoms are induced by enlargement, or hyperplasia, of the prostate gland. As the prostate increases in size, it impinges on free-flow of fluids through the male urethra. Concomitantly, the increased noradrenergic innervation of the enlarged prostate leads to an increased adrenergic tone of the bladder neck and urethra, further restricting the flow of urine through the urethra.

The α_2 receptors appear to have diverged rather early from either β or α_1 receptors. The α_2 receptors have been broken down into 3 molecularly distinct subtypes termed α_2C2 , α_2C4 , and α_2C10 based on their chromosomal location. These subtypes appear to correspond to the pharmacologically defined α_2B , α_2C , and α_2A subtypes, respectively (Bylund et al., 1992). While all the receptors of the adrenergic type are recognized by epinephrine, they are pharmacologically distinct and are encoded by separate genes. These receptors are generally coupled to different second messenger pathways that are linked through G-proteins. Among the adrenergic receptors, β_1 and β_2 receptors activate the adenylate cyclase, α_2 receptors inhibit adenylate cyclase and α_1 receptors activate phospholipase C pathways, stimulating breakdown of polyphosphoinositides (Chung, F. Z. et al., J. Biol. Chem., 263:4052 (1988)). α_1 and α_2 adrenergic receptors differ in their cell activity for drugs.

Issued US patent that disclose the utility of members of this family of proteins include, but are not limited to, 6,063,785 Phthalimido arylpiperazines useful in the treatment of benign prostatic hyperplasia; 6,060,492 Selective β_3 adrenergic agonists; 6,057,350 Alpha 1a adrenergic receptor antagonists; 6,046,192 Phenylethanolaminotetralincarboxamide derivatives; 6,046,183 Method of synergistic treatment for benign prostatic hyperplasia; 6,043,253 Fused piperidine substituted arylsulfonamides as β_3 -agonists; 6,043,224 Compositions and methods for treatment of neurological disorders and neurodegenerative diseases; 6,037,354 Alpha 1a adrenergic receptor antagonists; 6,034,106 Oxadiazole benzenesulfonamides as selective β_3 Agonist for the treatment of Diabetes and Obesity; 6,011,048 Thiazole benzenesulfonamides as β_3 agonists for treatment of diabetes and obesity; 6,008,361 5,994,506 Adrenergic receptor; 5,994,294 Nitrosated and nitrosylated α -adrenergic receptor antagonist compounds, compositions and their uses; 5,990,128 α_{1C} specific compounds to treat benign prostatic hyperplasia; 5,977,154 Selective β_3 adrenergic agonist; 5,977,115 Alpha 1a adrenergic receptor antagonists; 5,939,443 Selective β_3 adrenergic agonists; 5,932,538 Nitrosated and nitrosylated α -adrenergic receptor antagonist compounds, compositions and their uses; 5,922,722 Alpha 1a

adrenergic receptor antagonists 26 5,908,830 and 5,861,309 DNA encoding human alpha 1 adrenergic receptors.

Purinergic GPCRs

5 Purinoceptor P2Y1

P2 purinoceptors have been broadly classified as P2X receptors which are ATP-gated channels; P2Y receptors, a family of G protein-coupled receptors, and P2Z receptors, which mediate nonselective pores in mast cells. Numerous subtypes have been identified for each of the P2 receptor classes. P2Y receptors are characterized by their selective responsiveness towards ATP and its
10 analogs. Some respond also to UTP. Based on the recommendation for nomenclature of P2 purinoceptors, the P2Y purinoceptors were numbered in the order of cloning. P2Y1, P2Y2 and P2Y3 have been cloned from a variety of species. P2Y1 responds to both ADP and ATP. Analysis of P2Y receptor subtype expression in human bone and 2 osteoblastic cell lines by RT-PCR showed that all known human P2Y receptor subtypes were expressed: P2Y1, P2Y2, P2Y4, P2Y6, and P2Y7 (Maier et
15 al. 1997). In contrast, analysis of brain-derived cell lines suggested that a selective expression of P2Y receptor subtypes occurs in brain tissue.

Leon et al. generated P2Y1-null mice to define the physiologic role of the P2Y1 receptor (J. Clin. Invest. 104: 1731-1737(1999)). These mice were viable with no apparent abnormalities affecting their development, survival, reproduction, or morphology of platelets, and the platelet count in these
20 animals was identical to that of wildtype mice. However, platelets from P2Y1-deficient mice were unable to aggregate in response to usual concentrations of ADP and displayed impaired aggregation to other agonists, while high concentrations of ADP induced platelet aggregation without shape change. In addition, ADP-induced inhibition of adenylyl cyclase still occurred, demonstrating the existence of an ADP receptor distinct from P2Y1. P2Y1-null mice had no spontaneous bleeding tendency but were
25 resistant to thromboembolism induced by intravenous injection of ADP or collagen and adrenaline. Hence, the P2Y1 receptor plays an essential role in thrombotic states and represents a potential target for antithrombotic drugs. Somers et al. mapped the P2RY1 gene between flanking markers D3S1279 and D3S1280 at a position 173 to 174 cM from the most telomeric markers on the short arm of chromosome 3. (Genomics 44: 127-130 (1997)).

30 Purinoceptor P2Y2

The chloride ion secretory pathway that is defective in cystic fibrosis (CF) can be bypassed by an alternative pathway for chloride ion transport that is activated by extracellular nucleotides. Accordingly, the P2 receptor that mediates this effect is a therapeutic target for improving chloride

secretion in CF patients. Parr et al. reported the sequence and functional expression of a cDNA cloned from human airway epithelial cells that encodes a protein with properties of a P2Y nucleotide receptor. (Proc. Nat. Acad. Sci. 91: 3275-3279 (1994)) The human P2RY2 gene was mapped to chromosome 11q13.5-q14.1.

5 Purinoreceptor P2RY4

The P2RY4 receptor appears to be activated specifically by UTP and UDP, but not by ATP and ADP. Activation of this uridine nucleotide receptor resulted in increased inositol phosphate formation and calcium mobilization. The UNR gene is located on chromosome Xq13.

Purinoreceptor P2Y6

10 Somers et al. mapped the P2RY6 gene to 11q13.5, between polymorphic markers D11S1314 and D11S916, and P2RY2 maps within less than 4 cM of P2RY6. (Genomics 44: 127-130 (1997)) This was the first chromosomal clustering of this gene family to be described.

 Adenine and uridine nucleotides, in addition to their well established role in intracellular energy metabolism, phosphorylation, and nucleic acid synthesis, also are important extracellular
15 signaling molecules. P2Y metabotropic receptors are GPCRs that mediate the effects of extracellular nucleotides to regulate a wide variety of physiological processes. At least ten subfamilies of P2Y receptors have been identified. These receptor subfamilies differ greatly in their sequences and in their nucleotide agonist selectivities and efficacies.

 It has been demonstrated that the P2Y1 receptors are strongly expressed in the brain, but the
20 P2Y2, P2Y4 and P2Y6 receptors are also present. The localisation of one or more of these subtypes on neurons, on glia cells, on brain vasculature or on ventricle ependymal cells was found by in situ mRNA hybridisation and studies on those cells in culture. The P2Y1 receptors are prominent on neurons. The coupling of certain P2Y receptor subtypes to N-type Ca²⁺ channels or to particular K⁺ channels was also demonstrated.

25 It has also been demonstrated that several P2Y receptors mediate potent growth stimulatory effects on smooth muscle cells by stimulating intracellular pathways including Gq-proteins, protein kinase C and tyrosine phosphorylation, leading to increased immediate early gene expression, cell number, DNA and protein synthesis. It has been further demonstrated that P2Y regulation plays a mitogenic role in response to the development of atherosclerosis.

30 It has further been demonstrated that P2Y receptors play a critical role in cystic fibrosis. The volume and composition of the liquid that lines the airway surface is modulated by active transport of ions across the airway epithelium. This in turn is regulated both by autonomic agonists acting on basolateral receptors and by agonists acting on luminal receptors. Specifically, extracellular

nucleotides present in the airway surface liquid act on luminal P2Y receptors to control both Cl⁻ secretion and Na⁺ absorption. Since nucleotides are released in a regulated manner from airway epithelial cells, it is likely that their control over airway ion transport forms part of an autocrine regulatory system localised to the luminal surface of airway epithelia. In addition to this physiological role, P2Y receptor agonists have the potential to be of crucial benefit in the treatment of CF, a disorder of epithelial ion transport. The airways of people with CF have defective Cl⁻ secretion and abnormally high rates of Na⁺ absorption. Since P2Y receptor agonists can regulate both these ion transport pathways they have the potential to pharmacologically bypass the ion transport defects in CF.

10 Chemokine receptors:

The chemokines are structurally related proteins that act as chemoattractants and activators of lymphocytes and phagocytes. There are 2 separate families of chemokines differentiated by the location of the first 2 of 4 conserved cysteine residues. The alpha family is distinguished by the fact that the first 2 cysteines are separated by a single amino acid (CXC), while in the beta family the cysteines are adjacent (CC). The majority of the alpha chemokines, which includes IL8, target neutrophils, while the beta family members act largely upon monocytes. Members of the beta-chemokine family include macrophage inflammatory protein 1 alpha (MIP1-alpha), MIP1-beta, RANTES (regulated on activation, normal T expressed and secreted), MCP-1 (monocyte chemoattractant protein 1), MCP-2, MCP-3 and I-309. Receptors for chemokines have been cloned which have features of the G protein-coupled receptors.

Acute lung injury and the adult respiratory distress syndrome complicate many disease states. The mechanisms underlying this syndrome are unresolved, but the uniform pathologic features of adult respiratory distress syndrome involve sequestration of activated inflammatory cells within the lung, pulmonary microvascular injury, and leakage of intravascular fluid into the tissue spaces. In rodents, a model of these processes dependent on the initiation of acute pancreatitis is produced by overstimulation of pancreatic exocrine acinar cells with a cholecystokinin analog. Gerard et al. (1997) demonstrated that targeted disruption of the CCR1 receptor is associated with protection from pulmonary inflammation secondary to acute pancreatitis in the mouse. The protection from lung injury is associated with decreased levels of TNF-alpha in a temporal sequence indicating that the activation of the CCR1 receptor is an early event in the systemic inflammatory response syndrome.

Identification of the CC-chemokines RANTES, MIP1-alpha, and MIP1-beta as suppressor factors produced by CD8 cells that counter infection by certain HIV-1 strains facilitated the identification of 2 chemokine receptors, CXCR4 (or fusin) and CCR5, as cell surface co-receptors

with CD4 in HIV-1 infection. Additional receptors, CCR2 and CCR3, were also implicated as HIV-1 co-receptors on certain cell types. The findings in CCR5 and CXCR4 prompted a search for polymorphisms in other chemokine receptor genes that mediate disease progression. Smith et al. (1997) identified a val64-to-ile polymorphism (64I) in the first transmembrane region of CCR2, at an allele frequency of 10 to 15% among Caucasians and African Americans. Studies of 2 cohorts of AIDS patients showed that the CCR2-64I allele exerted no influence on the incidence of HIV-1 infection, but that HIV-1 infected persons carrying the 64I allele progressed to AIDS 2 to 4 years later than persons homozygous for the more common allele. Rapid progression of less than 3 years from HIV-1 exposure to onset of AIDS symptoms in an estimated 38 to 45% of AIDS patients could be attributed to their wildtype status at one or the other of these loci, whereas the survival of 28 to 29% of long-term survivors, who avoided AIDS for 16 years or more, could be explained by a mutant genotype for CCR2 or CCR5.

Chemokines are proinflammatory cytokines that function in leukocyte chemoattraction and activation. In addition to their function in viral disease, as describe above, chemokines have been implicated in the pathogenesis of atherosclerosis. Expression of the CC chemokine MCP1 is upregulated in human atherosclerotic plaques, in arteries of primates on a hypercholesterolemic diet, and in vascular endothelial and smooth muscle cells exposed to minimally modified lipids. To determine whether MCP1 is causally related to the development of atherosclerosis, Boring et al. (1998) generated mice that lacked CCR2, the receptor for MCP1, and crossed them with mice null for the apolipoprotein E gene (APOE), which develop severe atherosclerosis. They found that the selective absence of CCR2 decreased lesion formation markedly in apoE ^{-/-} mice but had no effect on plasma lipid or lipoprotein concentrations. These data revealed a role for MCP1 in the development of early atherosclerotic lesions and suggested that upregulation of this chemokine by minimally oxidized lipids is an important link between hyperlipidemia and fatty streak formation.

For a review of chemokine receptor and seven transmembrane G protein coupled chemokine receptor-like protein, see the references of Gerard *et al.*, *J. Clin. Invest.* 100: 2022-2027, 1997, Smith *et al.*, *Science* 277: 959-965, 1997, Boring *et al.*, *Nature* 394: 894-897, 1998.

GPCRs, particularly members of the chemokine receptor subfamily of the present invention which has seven transmembrane domains, are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown GPCRs. The present invention advances the state of the art by providing a previously unidentified human GPCR.

SUMMARY OF THE INVENTION

The present invention is based in part on the identification of nucleic acid sequences that encode amino acid sequences of human GPCR peptides and proteins that are related to the chemokine receptor subfamily, allelic variants thereof and other mammalian orthologs thereof.

5 These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and serve as targets for the development of human therapeutic agents.

The proteins of the present inventions are GPCRs that participate in signaling pathways mediated by the chemokine receptor subfamily in cells that express these proteins. Experimental data
10 as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. As used herein, a "signaling pathway" refers to the modulation (e.g., stimulation or inhibition) of a cellular function/activity upon the binding of a ligand to the GPCR protein. Examples of such functions include mobilization of intracellular molecules that participate in a signal transduction pathway, e.g., phosphatidylinositol 4,5-bisphosphate (PIP₂), inositol 1,4,5-triphosphate
15 (IP₃) and adenylate cyclase; polarization of the plasma membrane; production or secretion of molecules; alteration in the structure of a cellular component; cell proliferation, e.g., synthesis of DNA; cell migration; cell differentiation; and cell survival

The response mediated by the receptor protein depends on the type of cell it is expressed on. Some information regarding the types of cells that express other members of the subfamily of GPCRs
20 of the present invention is already known in the art (see references cited in Background and information regarding closest homologous protein provided in Figure 2; Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being.
) . For example, in some cells, binding of a ligand to the receptor protein may stimulate an activity such as release of compounds, gating of a channel, cellular adhesion, migration, differentiation, etc., through
25 phosphatidylinositol or cyclic AMP metabolism and turnover while in other cells, the binding of the ligand will produce a different result. Regardless of the cellular activity/response modulated by the particular GPCR of the present invention, a skilled artisan will clearly know that the receptor protein is a GPCR and interacts with G proteins to produce one or more secondary signals, in a variety of intracellular signal transduction pathways, e.g., through phosphatidylinositol or cyclic AMP
30 metabolism and turnover, in a cell thus participating in a biological process in the cells or tissues that express the GPCR. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being.

As used herein, "phosphatidylinositol turnover and metabolism" refers to the molecules involved in the turnover and metabolism of phosphatidylinositol 4,5-bisphosphate (PIP₂) as well as to the activities of these molecules. PIP₂ is a phospholipid found in the cytosolic leaflet of the plasma membrane. Binding of ligand to the receptor activates, in some cells, the plasma-membrane enzyme phospholipase C that in turn can hydrolyze PIP₂ to produce 1,2-diacylglycerol (DAG) and inositol 1,4,5-triphosphate (IP₃). Once formed IP₃ can diffuse to the endoplasmic reticulum surface where it can bind an IP₃ receptor, e.g., a calcium channel protein containing an IP₃ binding site. IP₃ binding can induce opening of the channel, allowing calcium ions to be released into the cytoplasm. IP₃ can also be phosphorylated by a specific kinase to form inositol 1,3,4,5-tetraphosphate (IP₄), a molecule that can cause calcium entry into the cytoplasm from the extracellular medium. IP₃ and IP₄ can subsequently be hydrolyzed very rapidly to the inactive products inositol 1,4-bisphosphate (IP₂) and inositol 1,3,4-triphosphate, respectively. These inactive products can be recycled by the cell to synthesize PIP₂. The other second messenger produced by the hydrolysis of PIP₂, namely 1,2-diacylglycerol (DAG), remains in the cell membrane where it can serve to activate the enzyme protein kinase C. Protein kinase C is usually found soluble in the cytoplasm of the cell, but upon an increase in the intracellular calcium concentration, this enzyme can move to the plasma membrane where it can be activated by DAG. The activation of protein kinase C in different cells results in various cellular responses such as the phosphorylation of glycogen synthase, or the phosphorylation of various transcription factors, e.g., NF-kB. The language "phosphatidylinositol activity", as used herein, refers to an activity of PIP₂ or one of its metabolites.

Another signaling pathway in which the receptor may participate is the cAMP turnover pathway. As used herein, "cyclic AMP turnover and metabolism" refers to the molecules involved in the turnover and metabolism of cyclic AMP (cAMP) as well as to the activities of these molecules. Cyclic AMP is a second messenger produced in response to ligand-induced stimulation of certain G protein coupled receptors. In the cAMP signaling pathway, binding of a ligand to a GPCR can lead to the activation of the enzyme adenylyl cyclase, which catalyzes the synthesis of cAMP. The newly synthesized cAMP can in turn activate a cAMP-dependent protein kinase. This activated kinase can phosphorylate a voltage-gated potassium channel protein, or an associated protein, and lead to the inability of the potassium channel to open during an action potential. The inability of the potassium channel to open results in a decrease in the outward flow of potassium, which normally repolarizes the membrane of a neuron, leading to prolonged membrane depolarization.

By targeting an agent to modulate a GPCR, the signaling activity and biological process mediated by the receptor can be agonized or antagonized in specific cells and tissues. Experimental

data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. Such agonism and antagonism serves as a basis for modulating a biological activity in a therapeutic context (mammalian therapy) or toxic context (anti-cell therapy, e.g. anti-cancer agent).

5

DESCRIPTION OF THE FIGURE SHEETS

FIGURE 1 provides the nucleotide sequence of a cDNA molecule or transcript sequence that encodes the GPCR of the present invention. (SEQ ID NO:1) In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of inventions based on this molecular sequence. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being.

FIGURE 2 provides the predicted amino acid sequence of the kinase of the present invention. (SEQ ID NO:2) In addition structure and functional information such as protein family, function, and modification sites is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence.

FIGURE 3 provides genomic sequences that span the gene encoding the GPCR protein of the present invention. (SEQ ID NO:3) In addition structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence.

DETAILED DESCRIPTION OF THE INVENTION

General Description

The present invention is based on the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a GPCR protein or part of a GPCR protein, that are related to the chemokine receptor subfamily. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human GPCR peptides and proteins that are related to the chemokine receptor subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these GPCR peptides and proteins, nucleic

acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the GPCR of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known GPCR proteins of the chemokine receptor subfamily and the expression pattern observed. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known chemokine family or subfamily of GPCR proteins.

Specific Embodiments

Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the GPCR family of proteins and are related to the chemokine receptor subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figure 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the GPCR peptides of the present invention, GPCR peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprise the amino acid sequences of the GPCR peptides disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be

based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

In some uses, "substantially free of cellular material" includes preparations of the peptide
5 having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes
10 preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the GPCR peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5%
15 chemical precursors or other chemicals.

The isolated GPCR peptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. For example, a nucleic acid molecule encoding the
20 GPCR peptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

Accordingly, the present invention provides proteins that consist of the amino acid sequences
25 provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence is the final amino acid sequence of the protein.

30 The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about

1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the GPCR peptides of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The GPCR peptides of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a GPCR peptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the GPCR peptide. "Operatively linked" indicates that the GPCR peptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the GPCR peptide.

In some uses, the fusion protein does not affect the activity of the GPCR peptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant GPCR peptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a

fusion moiety (e.g., a GST protein). A GPCR peptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the GPCR peptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the GPCR peptides of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence*

Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the GPCR peptides of the present invention as well as being encoded by the same genetic locus as the GPCR peptide provided herein. Mapping position in Figure 3 shows that the GPCR of the present invention is encoded by a gene on chromosome 12 near markers SHGC-64103 (LOD scores of 6.12), SHGC-56772 (LOD scores of 6.12) and SHGC-34758 (LOD scores of 6.06).

Allelic variants of a GPCR peptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the GPCR peptide as well as being encoded by the same genetic locus as the GPCR peptide provided herein. Genetic locus

can readily be determined based on the genomic information provided in Figure 3, such as the genomic sequence mapped to the reference human. Mapping position in Figure 3 shows that the GPCR of the present invention is encoded by a gene on chromosome 12 near markers SHGC-64103 (LOD scores of 6.12), SHGC-56772 (LOD scores of 6.12) and SHGC-34758 (LOD scores of 6.06). As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a GPCR peptide encoding nucleic acid molecule under stringent conditions as more fully described below.

10

Paralogs of a GPCR peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the GPCR peptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a GPCR peptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a GPCR peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the GPCR peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a GPCR peptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the GPCR peptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the GPCR peptide. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a GPCR peptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance concerning which

amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant GPCR peptides can be fully functional or can lack function in one or more activities, e.g. ability to bind ligand, ability to bind G-protein, ability to mediate signaling, etc. Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Figure 2 provides the result of protein analysis that identifies critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as ligand/effector molecule binding or in assays such as an *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

The present invention further provides fragments of the GPCR peptides, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a GPCR peptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the GPCR peptide or could be chosen for the ability to perform a function, e.g. ability to bind ligand or effector molecule or act as an immunogen. Particularly important fragments are biologically active fragments, peptides which are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the GPCR peptide, e.g., active site, a G-protein binding site, a transmembrane domain or a ligand-binding domain. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well-known and readily

available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in GPCR peptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well-known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the GPCR peptides of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature GPCR peptide is fused with another compound, such as a compound to increase the half-life of the GPCR peptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature GPCR peptide, such as a leader or secretory sequence or a sequence for purification of the mature GPCR peptide or a pro-protein sequence.

Protein/Peptide Uses

The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures and Back Ground Section; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to
5 quantitatively determine levels of the protein (or its binding partner or receptor) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the binding partner so as to develop a system
10 to identify inhibitors of the binding interaction. Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as commercial products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and
15 "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, GPCRs isolated from humans and their human/mammalian orthologs serve as targets for identifying agents for use in mammalian
20 therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the GPCR. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being. Approximately 70% of all pharmaceutical agents modulate the activity of
25 a GPCR. A combination of the invertebrate and mammalian ortholog can be used in selective screening methods to find agents specific for invertebrates. The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1. Experimental data as provided in Figure 1 indicates expression in the brain,
30 heart, lung, uterus, placenta and thyroid of human being. Such uses can readily be determined using the information provided herein, that known in the art and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to GPCRs that are related to members of the chemokine receptor subfamily. Such assays involve any of the known

GPCR functions or activities or properties useful for diagnosis and treatment of GPCR-related conditions that are specific for the subfamily of GPCRs that the one of the present invention belongs to, particularly in cells and tissues that express this receptor. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc.

5 Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being.

The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the receptor protein, as a biopsy or expanded in cell culture. Experimental data as provided in Figure 1 indicates
10 expression in the brain, heart, lung, uterus, placenta and thyroid of human being. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the receptor protein.

The polypeptides can be used to identify compounds that modulate receptor activity of the protein in its natural state, or an altered form that causes a specific disease or pathology associated with the receptor. Both the GPCRs of the present invention and appropriate variants and fragments can be
15 used in high-throughput screens to assay candidate compounds for the ability to bind to the receptor. These compounds can be further screened against a functional receptor to determine the effect of the compound on the receptor activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the receptor to a desired degree.

20 Further, the proteins of the present invention can be used to screen a compound for the ability to stimulate or inhibit interaction between the receptor protein and a molecule that normally interacts with the receptor protein, e.g. a ligand or a component of the signal pathway that the receptor protein normally interacts (for example, a G-protein or other interactor involved in cAMP or phosphatidylinositol turnover and/or adenylate cyclase, or phospholipase C activation). Such assays
25 typically include the steps of combining the receptor protein with a candidate compound under conditions that allow the receptor protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the receptor protein and the target, such as any of the associated effects of signal transduction such as G-protein phosphorylation, cAMP or phosphatidylinositol
30 turnover, and adenylate cyclase or phospholipase C activation.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of

random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules
5 obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the receptor that competes for ligand binding. Other candidate compounds include mutant receptors or appropriate fragments containing mutations that affect receptor function and thus compete for ligand. Accordingly, a fragment that competes for ligand, for example with a higher affinity, or a fragment that binds ligand but does not
10 allow release, is encompassed by the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) receptor activity. The assays typically involve an assay of events in the signal transduction pathway that indicate receptor activity. Thus, a cellular process such as proliferation, the expression of genes that are up- or down-regulated in response to the receptor protein dependent signal
15 cascade, can be assayed. In one embodiment, the regulatory region of such genes can be operably linked to a marker that is easily detectable, such as luciferase.

Any of the biological or biochemical functions mediated by the receptor can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and
20 other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological function of a cell or tissues that expresses the receptor can be assayed. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid
25 of human being.

Binding and/or activating compounds can also be screened by using chimeric receptor proteins in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by
30 heterologous domains or subregions. For example, a G-protein-binding region can be used that interacts with a different G-protein than that which is recognized by the native receptor. Accordingly, a different set of signal transduction components is available as an end-point assay for activation. Alternatively, the entire transmembrane portion or subregions (such as transmembrane segments or intracellular or extracellular loops) can be replaced with the entire transmembrane portion or

subregions specific to a host cell that is different from the host cell from which the amino terminal extracellular domain and/or the G-protein-binding region are derived. This allows for assays to be performed in other than the specific host cell from which the receptor is derived. Alternatively, the amino terminal extracellular domain (and/or other ligand-binding regions) could be replaced by a domain (and/or other binding region) binding a different ligand, thus, providing an assay for test compounds that interact with the heterologous amino terminal extracellular domain (or region) but still cause signal transduction. Finally, activation can be detected by a reporter gene containing an easily detectable coding region operably linked to a transcriptional regulatory sequence that is part of the native signal transduction pathway.

10 The proteins of the present invention are also useful in competition binding assays in methods designed to discover compounds that interact with the receptor. Thus, a compound is exposed to a receptor polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble receptor polypeptide is also added to the mixture. If the test compound interacts with the soluble receptor polypeptide, it decreases the amount of complex formed or activity from the receptor target. This type of assay is particularly useful in cases in which compounds are sought that
15 interact with specific regions of the receptor. Thus, the soluble polypeptide that competes with the target receptor region is designed to contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the receptor protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.
20

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., ³⁵S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated
25 by SDS-PAGE, and the level of receptor-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein
30

trapped in the wells by antibody conjugation. Preparations of a receptor-binding protein and a candidate compound are incubated in the receptor protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using
5 antibodies reactive with the receptor protein target molecule, or which are reactive with receptor protein and compete with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the GPCRs of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell
10 free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of receptor protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the receptor pathway, by treating cells or tissues that express the GPCR. Experimental data as provided in Figure 1 indicates expression in the brain,
15 heart, lung, uterus, placenta and thyroid of human being. These methods of treatment include the steps of administering a modulator of the GPCR's activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

In yet another aspect of the invention, the GPCR proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos et al. (1993)
20 *Cell* 72:223-232; Madura et al. (1993) *J. Biol. Chem.* 268:12046-12054; Bartel et al. (1993) *Biotechniques* 14:920-924; Iwabuchi et al. (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with the GPCR and are involved in GPCR activity. Such GPCR-binding proteins are also likely to be involved in the propagation of signals by the GPCR proteins or GPCR targets as, for example, downstream elements of a GPCR-mediated signaling pathway. Alternatively, such GPCR-binding proteins are likely to be GPCR
25 inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a GPCR protein is fused to a gene
30 encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a GPCR-dependent complex, the DNA-binding and activation domains of the transcription factor are

brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein
5 which interacts with the GPCR protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a GPCR modulating agent, an antisense GPCR nucleic acid molecule, a GPCR-specific
10 antibody, or a GPCR-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

15 The GPCR proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. The method involves contacting a biological sample with
20 a compound capable of interacting with the receptor protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as
25 well as tissues, cells and fluids present within a subject.

The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a
30 genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered receptor activity in cell-based or cell-free assay, alteration in ligand or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques

useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

In vitro techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the receptor protein in which one or more of the receptor functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other ligand-binding regions that are more or less active in ligand binding, and receptor activation. Accordingly, ligand dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data as provided in Figure 1

indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. Accordingly, methods for treatment include the use of the GPCR protein or fragments.

Antibodies

5 The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments thereof. As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so
10 long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

 As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen
15 challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')₂, and Fv fragments.

 Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, Antibodies, Cold Spring Harbor Press, (1989).

20 In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein
25 alignment methods and as presented in the Figures.

 Antibodies are preferably prepared from regions or discrete fragments of the GPCR proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or receptor/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence
30 alignment can be used to identify conserved and unique sequence fragments.

 An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as fragments correspond to regions that are

located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various
5 enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include
10 umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Antibody Uses

15 The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues
20 in an organism and over the course of normal development. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such
25 antibodies can be used to assess abnormal tissue distribution or abnormal expression during development or progression of a biological condition. Antibody detection of circulating fragments of the full length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function.
30 When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. If a disorder is characterized by a specific mutation in the

protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

5 The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

10 Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

15 The antibodies are also useful for tissue typing. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

20 The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the GPCR peptide to a binding partner such as a ligand. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information
25 relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in the sample with a standard; and
30 instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array. Arrays are described in detail below for nucleic acid arrays and similar methods have been developed for antibody arrays.

Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a GPCR peptide or protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the GPCR peptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3,

genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

5 The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid
10 molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprise several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

15 In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, human genomic sequences (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be
20 identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

25 The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is
30 the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the GPCR peptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the

sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form of DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides nucleic acid molecules that encode fragments of the peptides of the present invention as well as nucleic acid molecules that encode obvious variants of the GPCR proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such fragments are useful in controlling heterologous gene expression and in developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could be at least 30, 40, 50, 100, 250 or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid

corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under
5 stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence.
10 Such nucleic acid molecules can readily be identified as being able to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. Mapping position in Figure 3 shows that the GPCR of the present invention is encoded by a gene on chromosome 12 near markers SHGC-64103 (LOD scores of 6.12), SHGC-56772 (LOD scores of 6.12)
15 and SHGC-34758 (LOD scores of 6.06).

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions can be
20 such that sequences at least about 60%, at least about 70%, or at least about 80% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45C, followed by one or more washes in 0.2 X SSC,
25 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridization conditions are well known in the art.

Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical
30 intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as encompassing fragments disclosed prior to the present invention.

5 The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

 The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into
10 the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

 The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

 The nucleic acid molecules are also useful as probes for determining the chromosomal
15 positions of the nucleic acid molecules by means of *in situ* hybridization methods. Mapping position in Figure 3 shows that the GPCR of the present invention is encoded by a gene on chromosome 12 near markers SHGC-64103 (LOD scores of 6.12), SHGC-56772 (LOD scores of 6.12) and SHGC-34758 (LOD scores of 6.06).

 The nucleic acid molecules are also useful in making vectors containing the gene regulatory
20 regions of the nucleic acid molecules of the present invention.

 The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

 The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

25 The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

 The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

 The nucleic acid molecules are also useful as hybridization probes for determining the
30 presence, level, form and distribution of nucleic acid expression. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being. Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic

acid whose level is determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in GPCR protein expression relative to normal results.

5 *In vitro* techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a GPCR protein, such as by measuring a level of a receptor-encoding nucleic acid in a sample of cells
10 from a subject e.g., mRNA or genomic DNA, or determining if a receptor gene has been mutated. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being.

Nucleic acid expression assays are useful for drug screening to identify compounds that
15 modulate GPCR nucleic acid expression.

The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the GPCR gene, particularly biological and pathological processes that are mediated by the GPCR in cells and tissues that express it. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid
20 of human being. The method typically includes assaying the ability of the compound to modulate the expression of the GPCR nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired GPCR nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the GPCR nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

25 The assay for GPCR nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the GPCR protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

30 Thus, modulators of GPCR gene ~~expression~~ can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of GPCR mRNA in the presence of the candidate compound is compared to the level of expression of GPCR mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for

example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than
5 in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate GPCR nucleic acid expression, particularly to modulate activities within a cell or tissue that expresses the proteins. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are
10 expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or nucleic acid expression.

Alternatively, a modulator for GPCR nucleic acid expression can be a small molecule or drug
15 identified using the screening assays described herein as long as the drug or small molecule inhibits the GPCR nucleic acid expression in the cells and tissues that express the protein. Experimental data as provided in Figure 1 indicates expression in the brain, heart, lung, uterus, placenta and thyroid of human being.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating
20 compounds on the expression or activity of the GPCR gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased
25 administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in GPCR nucleic acid, and particularly in qualitative changes that lead to pathology. The nucleic acid
30 molecules can be used to detect mutations in GPCR genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally-occurring genetic mutations in the GPCR gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition,

modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the GPCR gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a GPCR protein.

5 Individuals carrying mutations in the GPCR gene can be detected at the nucleic acid level by a variety of techniques. Mapping position in Figure 3 shows that the GPCR of the present invention is encoded by a gene on chromosome 12 near markers SHGC-64103 (LOD scores of 6.12), SHGC-56772 (LOD scores of 6.12) and SHGC-34758 (LOD scores of 6.06). Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same
10 way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.*, *Nucleic Acids Res.*
15 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing
20 the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a GPCR gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

25 Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such
30 as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant GPCR gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g.,

PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship). Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the GPCR gene in an individual in order to select an appropriate compound or dosage regimen for treatment.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control GPCR gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of GPCR protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into GPCR protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of GPCR nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired GPCR nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the GPCR protein, such as ligand binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in GPCR gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired GPCR protein to treat the individual.

5 The invention also encompasses kits for detecting the presence of a GPCR nucleic acid in a biological sample. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in the brain, heart, lung etc. Specifically, a virtual northern blot shows expression the brain, heart, lung, uterus, placenta and thyroid of human being. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting GPCR
10 nucleic acid in a biological sample; means for determining the amount of GPCR nucleic acid in the sample; and means for comparing the amount of GPCR nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect GPCR protein mRNA or DNA.

15 Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or
20 oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee et al., PCT application W095/11995 (Chee et al.), Lockhart, D. J. et al. (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. et al. (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in
25 their entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown et. al., US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in
30 length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length

of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or detection kit, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler et al.) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray or detection kit. The biological samples may be

obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of the GPCR proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the GPCR gene of the present invention.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed herein.

Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified GPCR genes of the present invention can be routinely identified using the sequence information disclosed herein can be readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in procaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood,

however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, eg. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an

expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene* 69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in a host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

5 The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory*
10 *Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.*

 The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a
15 portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

 The invention also relates to recombinant host cells containing the vectors described herein.
20 Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

 The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic
25 lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989*).

 Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be
30 introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

5 Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides
10 selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

15 Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as GPCRs, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

Where the peptide is not secreted into the medium, which is typically the case with GPCRs, the
20 protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography,
25 lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

30

Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a GPCR protein or peptide that can be further purified to

produce desired amounts of GPCR protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the GPCR protein or GPCR protein fragments, such as those described above as well as other formats known in the art.

- 5 Thus, a recombinant host cell expressing a native GPCR protein is useful for assaying compounds that stimulate or inhibit GPCR protein function.

- Host cells are also useful for identifying GPCR protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant GPCR protein (for
10 example, stimulating or inhibiting function) which may not be indicated by their effect on the native GPCR protein.

- Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA
15 which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a GPCR protein and identifying and evaluating modulators of GPCR protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

- 20 A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the GPCR protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

- Any of the regulatory or other sequences useful in expression vectors can form part of the
25 transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the GPCR protein to particular cells.

- Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example,
30 in U.S. Patent Nos. 4,736,866 and 4,870,069, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder

animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

5 In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al. PNAS* 89:6232-6236 (1992). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al. Science* 251:1351-1355 (1991). If a
10 *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

15 Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from
20 an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are
25 useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect ligand binding, GPCR protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* GPCR protein function, including ligand interaction, the effect of specific mutant GPCR proteins on GPCR protein
30 function and ligand interaction, and the effect of chimeric GPCR proteins. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more GPCR protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the

invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for

5 carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

<110> Gilbert et al.

<120> ISOLATED HUMAN G-PROTEIN COUPLED
RECEPTORS, NUCLEIC ACID MOLECULES ENCODING HUMAN GPCR
PROTEINS, AND USES THEREOF

<130> CL000782PCT

<140> N/A

<141> 2001-03-27

<150> US 60/192,419

<151> 2000-03-27

<150> US 60/230,459

<151> 2000-09-06

<150> US 09/666,535

<151> 2000-09-20

<160> 4

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1041

<212> DNA

<213> Human

<400> 1

```

atgtacaacg ggtcgtgctg ccgcatcgag ggggacacca tctcccaggt gatgccgccc 60
ctgctcattg tggcccttgt gctgggcgca ctaggcaatg gggtcgccct gtgtggtttc 120
tgcttcacac tgaagacctg gaagcccagc actgtttacc ttttcaattt ggccgtgggt 180
gatttcctcc ttatgatctg cctgcctttt cggacagact attacctcag acgtagacac 240
tgggcttttg gggacattcc ctgcccagtg gggctcttca cgttggccat gaacagggcc 300
gggagcatcg tgttccttac ggtggtggct gcggacaggt atttcaaagt ggtccacccc 360
caccacgctg tgaacactat ctccaccggt gtggcgctg gcatcgctctg caccctgtgg 420
gccctggtca tccctgggaac agtgtatctt ttgctggaga accatctctg cgtgcaagag 480
acggccgtct cctgtgagag cttcatcatg gagtcggcca atggctggca tgacatcatg 540
ttccagctgg agttctttat gccctcggtc atcatcttat ttgtctcctt caagattggt 600
tggagcctga ggcggaggca gcagctggcc agacaggctc ggatgaagaa ggcgaccccg 660
ttcatcatgg tgggtgcaat tgtgttcac acatgctacc tgcccagcgt gtctgctaga 720
ctctatttcc tctggacggt gccctcgagt gcctgcgac cctctgtcca tggggccctg 780
cacataaccc tcagcttcac ctacatgaac agcatgctgg atcccctggt gtattatttt 840
tcaagcccct cctttcccaa attctacaac aagctcaaaa tctgcagtct gaaacccaag 900
cagccaggac actcaaaaac acaaggccg gaagagatgc caatttcgaa cctcggtcgc 960
aggagtgcga tcagtgtggc aaatagtttc caaagccagt ctgatgggca atgggatccc 1020
cacattgttg agtggcactg a

```

<210> 2

<211> 346

<212> PRT

<213> Human

<400> 2

```

Met Tyr Asn Gly Ser Cys Cys Arg Ile Glu Gly Asp Thr Ile Ser Gln
1           5           10           15
Val Met Pro Pro Leu Leu Ile Val Ala Phe Val Leu Gly Ala Leu Gly
20           25           30
Asn Gly Val Ala Leu Cys Gly Phe Cys Phe His Met Lys Thr Trp Lys
35           40           45

```

```

Pro Scr Thr Val Tyr Leu Phe Asn Leu Ala Val Ala Asp Phe Leu Leu
  50                      55                      60
Met Ile Cys Leu Pro Phe Arg Thr Asp Tyr Tyr Leu Arg Arg Arg His
  65                      70                      75                      80
Trp Ala Phe Gly Asp Ile Pro Cys Arg Val Gly Leu Phe Thr Leu Ala
                      85                      90                      95
Met Asn Arg Ala Gly Ser Ile Val Phe Leu Thr Val Val Ala Ala Asp
                      100                      105                      110
Arg Tyr Phe Lys Val Val His Pro His His Ala Val Asn Thr Ile Ser
                      115                      120                      125
Thr Arg Val Ala Ala Gly Ile Val Cys Thr Leu Trp Ala Leu Val Ile
                      130                      135                      140
Leu Gly Thr Val Tyr Leu Leu Leu Glu Asn His Leu Cys Val Gln Glu
  145                      150                      155                      160
Thr Ala Val Ser Cys Glu Ser Phe Ile Met Glu Ser Ala Asn Gly Trp
                      165                      170                      175
His Asp Ile Met Phe Gln Leu Glu Phe Phe Met Pro Leu Gly Ile Ile
                      180                      185                      190
Leu Phe Cys Ser Phe Lys Ile Val Trp Ser Leu Arg Arg Arg Gln Gln
                      195                      200                      205
Leu Ala Arg Gln Ala Arg Met Lys Lys Ala Thr Arg Phe Ile Met Val
                      210                      215                      220
Val Ala Ile Val Phe Ile Thr Cys Tyr Leu Pro Ser Val Ser Ala Arg
  225                      230                      235                      240
Leu Tyr Phe Leu Trp Thr Val Pro Ser Ser Ala Cys Asp Pro Ser Val
                      245                      250                      255
His Gly Ala Leu His Ile Thr Leu Ser Phe Thr Tyr Met Asn Ser Met
                      260                      265                      270
Leu Asp Pro Leu Val Tyr Tyr Phe Ser Ser Pro Ser Phe Pro Lys Phe
                      275                      280                      285
Tyr Asn Lys Leu Lys Ile Cys Ser Leu Lys Pro Lys Gln Pro Gly His
                      290                      295                      300
Ser Lys Thr Gln Arg Pro Glu Glu Met Pro Ile Ser Asn Leu Gly Arg
  305                      310                      315                      320
Arg Ser Cys Ile Ser Val Ala Asn Ser Phe Gln Ser Gln Ser Asp Gly
                      325                      330                      335
Gln Trp Asp Pro His Ile Val Glu Trp His
                      340                      345

```

<210> 3
 <211> 2334
 <212> DNA
 <213> Human

```

<400> 3
aatgaccac ttttgcaaaa ttgcatgcat ttccaagctt catccggctc . caggcttg 60
gcctctccca gaggcaggcg gcttgtgaga cgggctccag ag aaaggac ctccctgggt 120
ctctcatttc ctgggtgaag tttctcttct cgctgctgtg gcagcatcca acccacacac 180
acaggaccgc catcctgggt gatgaagtca gacacgcagc agctgggtga gtgctaacgc 240
tcagataagc atctgtgcca ttgtggggac tccctgggct gctctgcacc cggacacttg 300
ctctgtcccc gccatgtaca acgggtcgtg ctgccgcatac gagggggaca ccatctccca 360
gggtgatgcc cgctgtctca ttgtggcctt tgtgctgggc gcactaggca atggggctcg 420
cctgtgtggt ttctgcttcc acatgaagac ctggaagccc agcactgttt accttttcaa 480
tttggccgtg gctgatttcc tccttatgat ctgcctgcct tttcggacag actattacct 540
cagacgtaga cactgggctt ttggggacat tccctgccga gtggggctct tcacgttggc 600
catgaacagg gccgggagca tcgtgttcc taccggtggtg gctgcggaca ggtatttcaa 660
agtqqtccac cccaccacg cgggtgaacac tatctccacc cgggtggcgg ctggcatcgt 720
ctgcaccctg tgggccctgg tcctcctggg aacagtgtat cttttgctgg agaaccatct 780
ctgctgtcaa gagacggccg tctcctgtga gagcttcatac atggagtcgg ccaatggctg 840
gcatgacatc atgttccagc tggagtctt tatgccctc ggcatcatct tattttgctc 900
cttcaagatt gtttggagcc tgaggcggag gcagcagctg gccagacagg ctcgatgaa 960
gaaggcgacc cgttccatca tgggtgtggc aattgtgttc atcacatgct acctgcccag 1020

```

```

cgtgtctgct agactctatt tcctctggac ggtgccctcg agtgccctgc atccctctgt 1080
ccatggggcc ctgcacataa ccttcagctt cacctacalg aacagcatgc tggatcccct 1140
ggtgtattat ttttcaagcc cctcctttcc caaatcttac aacaagctca aaatctgcag 1200
tctgaaaccc aagcagccag gacactcaaa aacacaaagg ccggaagaga tgccaatttc 1260
gaacctcgtt cgcaggagtt gcatcagtg ggcaaatagt ttccaaagcc agtctgatgg 1320
gcaatgggat cccacattg ttgagtggca ctgaacaagc agaccaacaa cactgaggaa 1380
gatagagtgg tgacttagaa ttaactcgtg ctaaggggtc gggggctttg aaaatgccac 1440
ccccctttct tattgcaaga cggcttctcg cacatgaact gcctccttct cattctgtcg 1500
gaaatgaaat tcacacaact ataccttttg gggaggttcc agttgattga agtgagttgg 1560
ctgcattttc ttatctgac acaatggcag gggacagaat gtgcatggag tggagcatgt 1620
gtgtgttggg aggggggcta ggaactgcac agcccttggt taatttctgt tgtttgtttt 1680
tgttttgaga cagagtctca ctctgtgtcc caggctggag tgcagtggca cagtctcggc 1740
tcactgcaac ctctgcctcc cgggttcaag caattctcct gcctcagcct cccgagtagc 1800
tggtattaga ggcgccagcc aacacaccgc gctaattttt gtatttttag tagagacagg 1860
gttttgccat gttggccagg ctggtctcga gctcctgacc tcaggtgatc cgctgcctt 1920
ggcctcccaa agtggtgagg tcacaggcgt gagccaccgt gcccggcctc cctgtgttca 1980
ttttaaatgg ctaagtaa at gggtatatgt gtttgaatgg ggcagtgtca ctctcttagg 2040
ggctatgggg cagttagcag catttcttat cctclgacct taaatcattc cttatctcag 2100
aaaacagaaa ccgggctcag tcaatcaatg ctttatttca ggccgaatga ggctcttttag 2160
attgggatct attgatctat caattttcat ctttacattt ctttgtacat ctgtacattt 2220
tgtccaaatg tacatctgta cgtctgtcat cattgtgact tcctggtagc ccaagaagaa 2280
caacaacaaa acaatctgct ctgaccttct tcaaatcttt gtatttcaaa gaag 2334

```

<210> 4
 <211> 339
 <212> PRT
 <213> Human

```

<400> 4
Asn Cys Cys Val Phe Arg Asp Asp Phe Ile Ala Lys Val Leu Pro Pro
 1          5          10          15
Val Leu Gly Leu Glu Phe Ile Phe Gly Leu Leu Gly Asn Gly Leu Ala
 20          25          30
Leu Trp Ile Phe Cys Phe His Leu Lys Ser Trp Lys Ser Arg Ile
 35          40          45
Phe Leu Phe Asn Leu Ala Val Ala Asp Phe Leu Leu Ile Ile Cys Leu
 50          55          60
Pro Phe Val Met Asp Tyr Tyr Val Arg Arg Ser Asp Trp Asn Phe Gly
 65          70          75          80
Asp Ile Pro Cys Arg Leu Val Leu Phe Met Phe Ala Met Asn Arg Gln
 85          90          95
Gly Ser Ile Ile Phe Leu Thr Val Val Ala Val Asp Arg Tyr Phe Arg
100          105          110
Val Val His Pro His His Ala Leu Asn Lys Ile Ser Asn Trp Thr Ala
115          120          125
Ala Ile Ile Ser Cys Leu Leu Trp Gly Ile Thr Val Gly Leu Thr Val
130          135          140
His Leu Leu Lys Lys Lys Leu Leu Ile Gln Asn Gly Pro Ala Asn Val
145          150          155          160
Cys Ile Ser Phe Ser Ile Cys His Thr Phe Arg Trp His Glu Ala Met
165          170          175
Phe Leu Leu Glu Phe Leu Leu Pro Leu Gly Ile Ile Leu Phe Cys Ser
180          185          190
Ala Arg Ile Ile Trp Ser Leu Arg Gln Arg Gln Met Asp Arg His Ala
195          200          205
Lys Ile Lys Arg Ala Ile Thr Phe Ile Met Val Val Ala Ile Val Phe
210          215          220
Val Ile Cys Phe Leu Pro Ser Val Val Val Arg Ile Arg Ile Phe Trp
225          230          235          240
Leu Leu His Thr Ser Gly Thr Gln Asn Cys Glu Val Tyr Arg Ser Val
245          250          255
Asp Leu Ala Phe Phe Ile Thr Leu Ser Phe Thr Tyr Met Asn Ser Met
260          265          270

```


Leu Asp Pro Val Val Tyr Tyr Phe Ser Ser Pro Ser Phe Pro Asn Phe
275 280 285
Phe Ser Thr Leu Ile Asn Arg Cys Leu Gln Arg Lys Met Thr Gly Glu
290 295 300
Pro Asp Asn Asn Arg Ser Thr Ser Val Glu Leu Thr Gly Asp Pro Asn
305 310 315 320
Lys Thr Arg Gly Ala Pro Glu Ala Leu Met Ala Asn Ser Gly Glu Pro
325 330 335
Trp Ser Pro

Claims

That which is claimed is:

1. An isolated peptide consisting of an amino acid sequence selected from the group consisting of:
 - (a) an amino acid sequence shown in SEQ ID NO:2;
 - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
 - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic); and
 - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
2. An isolated peptide comprising an amino acid sequence selected from the group consisting of:
 - (a) an amino acid sequence shown in SEQ ID NO:2;
 - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
 - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic); and
 - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
3. An isolated antibody that selectively binds to a peptide of claim 2.

4. An isolated nucleic acid molecule consisting of a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

5. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

6. A gene chip comprising a nucleic acid molecule of claim 5.

7. A transgenic non-human animal comprising a nucleic acid molecule of claim 5.
8. A nucleic acid vector comprising a nucleic acid molecule of claim 5.
9. A host cell containing the vector of claim 8.
10. A method for producing any of the peptides of claim 1 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
11. A method for producing any of the peptides of claim 2 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
12. A method for detecting the presence of any of the peptides of claim 2 in a sample, said method comprising contacting said sample with a detection agent that specifically allows detection of the presence of the peptide in the sample and then detecting the presence of the peptide.
13. A method for detecting the presence of a nucleic acid molecule of claim 5 in a sample, said method comprising contacting the sample with an oligonucleotide that hybridizes to said nucleic acid molecule under stringent conditions and determining whether the oligonucleotide binds to said nucleic acid molecule in the sample.
14. A method for identifying a modulator of a peptide of claim 2, said method comprising contacting said peptide with an agent and determining if said agent has modulated the function or activity of said peptide.
15. The method of claim 14, wherein said agent is administered to a host cell comprising an expression vector that expresses said peptide.
16. A method for identifying an agent that binds to any of the peptides of claim 2, said method comprising contacting the peptide with an agent and assaying the contacted mixture to determine whether a complex is formed with the agent bound to the peptide.

17. A pharmaceutical composition comprising an agent identified by the method of claim 16 and a pharmaceutically acceptable carrier therefor.
18. A method for treating a disease or condition mediated by a human proteases, said method comprising administering to a patient a pharmaceutically effective amount of an agent identified by the method of claim 16.
19. A method for identifying a modulator of the expression of a peptide of claim 2, said method comprising contacting a cell expressing said peptide with an agent, and determining if said agent has modulated the expression of said peptide.
20. An isolated human protease peptide having an amino acid sequence that shares at least 70% homology with an amino acid sequence shown in SEQ ID NO:2.
21. A peptide according to claim 20 that shares at least 90 percent homology with an amino acid sequence shown in SEQ ID NO:2.
22. An isolated nucleic acid molecule encoding a human protease peptide, said nucleic acid molecule sharing at least 80 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic).
23. A nucleic acid molecule according to claim 22 that shares at least 90 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic).

1/5

FIGURE 1

```

1  ATGTACAACG GGTCTGCTG. CCGCATCGAG GGGGACACCA TCTCCCAGGT
51  GATGCCGCCG CTGCTCATTG TGGCCTTTGT GCTGGGCGCA CTAGGCAATG
101 GGGTCGCCCT GTGTGGTTTC TGCTTCCACA TGAAGACCTG GAAGCCCAGC
151 ACTGTTTACC TTTTCAATTG GGCCGTGGCT GATTTCCTCC TTATGATCTG
201 CCTGCCTTTT CGGACAGACT ATTACCTCAG ACGTAGACAC TGGGCTTTTG
251 GGGACATTCC CTGCCGAGTG GGGCTCTTCA CGTTGGCCAT GAACAGGGCC
301 GGGAGCATCG TGTTCTTAC GGTGGTGGCT GCGGACAGGT ATTTCAAAGT
351 GGTCCACCCC CACCACGCGG TGAACACTAT CTCCACCCGG GTGGCGGCTG
401 GCATCGTCTG CACCCTGTGG GCCCTGGTCA TCCTGGGAAC AGTGTATCTT
451 TTGCTGGAGA ACCATCTCTG CGTGCAAGAG ACGGCCGTCT CCTGTGAGAG
501 CTTTCATCATG GAGTCGGCCA ATGGCTGGCA TGACATCATG TTCCAGCTGG
551 AGTTCTTTAT GCCCCTCGGC ATCATCTTAT TTTGCTCCTT CAAGATTGTT
601 TGGAGCCTGA GCGGAGGCA GCAGCTGGCC AGACAGGCTC GGATGAAGAA
651 GCGGACCCGG TTCATCATGG TGGTGGCAAT TGTGTTCAAT ACATGCTACC
701 TGCCAGCGT GTCTGCTAGA CTCTATTTCC TCTGGACGGT GCCCTCGAGT
751 GCCTGCGATC CCTCTGTCCA TGGGGCCCTG CACATAACCC TCAGCTTCAC
801 CTACATGAAC AGCATGCTGG ATCCCCTGGT GTATTATTTT TCAAGCCCCT
851 CCTTTCCCAA ATTCTACAAC AAGCTCAAAA TCTGCAGTCT GAAACCCAAG
901 CAGCCAGGAC ACTCAAAAAC ACAAAGGCCG GAAGAGATGC CAATTTGAA
951 CCTCGGTCGC AGGAGTTGCA TCAGTGTGGC AAATAGTTTC CAAAGCCAGT
1001 CTGATGGGCA ATGGGATCCC CACATTGTTG AGTGGCACTG A (SEQ ID NO:1)

```

FEATURES:

Start: 1

Stop: 1039

HOMOLOGOUS PROTEIN:Top 10 BLAST Hits:

gi 5174461 ref NP_006009.1 putative chemokine receptor; GTP-b...	344	7e-94
gi 4885317 ref NP_005290.1 G protein-coupled receptor 31 >gi ...	175	6e-43
gi 6716509 gb AAF26668.1 AF140708_1 (AF140708) G protein couple...	172	4e-42
gi 464327 sp P34996 P2YR_CHICK P2Y PURINOCEPTOR 1 (ATP RECEPTOR...	138	6e-32
gi 1352693 sp P49652 P2YR_MELGA P2Y PURINOCEPTOR 1 (ATP RECEPTO...	138	6e-32
gi 1352691 sp P48042 P2YR_BOVIN P2Y PURINOCEPTOR 1 (ATP RECEPTO...	138	1e-31
gi 4505557 ref NP_002554.1 purinergic receptor P2Y, G-protein...	135	5e-31
gi 1352695 sp P49651 P2YR_RAT P2Y PURINOCEPTOR 1 (ATP RECEPTOR)...	135	5e-31
gi 6679193 ref NP_032798.1 purinergic receptor P2Y, G-protein...	131	8e-30
gi 4885311 ref NP_005287.1 G protein-coupled receptor 23 >gi ...	130	2e-29

EXPRESSION INFORMATION FOR MODULATORY USE (from cDNA panel screening):

Human Brain
 Human Heart
 Human Lung
 Human Uterus
 Human Placenta
 Human Thyroid

2/5

FIGURE 2, page 1 of 2

```

1  MYNGSCCRIE GDTISQVMPP LLIVAFVLGA LGNGVALCGF CFHMKTWKPS
51 TVYLFNLAVA DFLLMICLPF RTDYLLRRRH WAFGDIPCRV GLFTLAMNRA
101 GSIVFLTVVA ADRYFKVVHP HHAVNTISTR VAAGIVCTLW ALVILGTVYL
151 LLENHLCVQE TAVSCSFIM ESANGWHDIM FQLEFFMPLG IILFCSFKIV
201 WSLRRRQQLA RQARMKKATR FIMVVAIVFI TCYLPSVSAR LYFLWTVPS
251 ACDPSVHGAL HITLSFTYMN SMLDPLVYFF SSPSPKPFYN KLIKICSLKPK
301 QPGHSTQRP EEMPISNLGR RSCISVANSE QSDGQWDP HIVEWH (SEQ ID NO:2)

```

FEATURES:**Functional domains and key regions:**

[1] PDOC00001 PS00001 ASN_GLYCOSYLATION

N-glycosylation site

3-6 NGSC

[2] PDOC00004 PS00004 CAMP_PHOSPHO_SITE

cAMP- and cGMP-dependent protein kinase phosphorylation site

216-219 KKAT

[3] PDOC00005 PS00005 PKC_PHOSPHO_SITE

Protein kinase C phosphorylation site

Number of matches: 7

```

1 46-48 TWK
2 128-130 STR
3 196-198 SFK
4 202-204 SLR
5 238-240 SAR
6 296-298 SLK
7 307-309 TQR

```

[4] PDOC00006 PS00006 CK2_PHOSPHO_SITE

Casein kinase II phosphorylation site

Number of matches: 3

```

1 250-253 SACD
2 271-274 SMLD
3 332-335 QSD

```

[5] PDOC00008 PS00008 MYRISTYL

N-myristoylation site

Number of matches: 4

```

1 29-34 GALGNG
2 32-37 GNGVAL
3 34-39 GVALCG
4 134-139 GIVCTL

```

[6] PDOC00009 PS00009 AMIDATION

Amidation site

318-321 LGRR

[7] PDOC00210 PS00237 G_PROTEIN_RECEPTOR

G-protein coupled receptors signature

101-117 GSIVFLTVVAADRYFKV

Membrane spanning structure and domains:

Helix	Begin	End	Score	Certainty
1	13	33	1.800	Certain
2	50	70	1.312	Certain
3	90	110	0.906	Putative
4	131	151	2.097	Certain
5	183	203	1.539	Certain
6	219	239	1.884	Certain
7	254	274	0.950	Putative

3/5

FIGURE 2, page 2 of 2

BLAST Alignment to Top Hit:

```

>gi|5174461|ref|NP_006009.1| putative chemokine receptor;
      GTP-binding protein >gi|1346295|sp|P49019|HM74_HUMAN
      PROBABLE G PROTEIN-COUPLED RECEPTOR HM74
>gi|2135386|pir||I69202 HM74 - human
>gi|219867|dbj|BAA01721.1| (D10923) HM74 [Homo sapiens]
      Length = 387

Score = 344 bits (873), Expect = 9e-94
Identities = 178/341 (52%), Positives = 227/341 (66%), Gaps = 7/341 (2%)

Query: 5   SCCRIEGDTISQVMPPLLIVAFVLGALNGVALCGFCFHMKTWKPSTVYLFNLAVADFL 64
          +CC   D I++V+PP+L + F+ G LGNG+AL FCFH+K+WK S ++LFNLAVADFL
Sbjct: 17  NCCVFRDDFIKVLPPVLGLEFIFGLLGNGLALWIFCFHLKSWKSSRIFLFNLAVADFL 76

Query: 65  MICLPFRTDYLLRRRHWAFGDIPCRVGLFTLAMNRAGSIVFLTVAADRYFKVVPHHAV 124
          +ICLPF DYY+RR W FGDIPCR+ LF AMNR GSI+FLTVA DRYF+VVPHHA+
Sbjct: 77  IICLPFMDYVRRSDWNFGDIPCRVLVLFMFAMNRQGSIIFLTVAVDYFRVVPHHAL 136

Query: 125 NTISTRVAAGIVCTLWALVILGTVYLLLENHLCVQETAVSCSFIMESANGWHDIMFQLE 184
          N IS  AA I C LW + + TV+LL + L  A C SF +  WH+ MF LE
Sbjct: 137 NKISNWTAAIISCLLWGITVGLTVHLLKKKLLIQNGPANVCISFSICHTFRWHEAMFLE 196

Query: 185 FEMPLGIILFCSFKIVWSLRRRQQLARQARMKKATRFIMVVAIVFITCYLPSVSARLYFL 244
          F +PLGIILFCS +I+WSLR+R Q+ R A++K+A FIMVVAIVE+ C+LPSV R+
Sbjct: 197 FLLPLGIILFCSARIIWSLRQR-QMDRHAKIKRAITFIMVVAIVFVICFLPSVVVRIRIF 255

Query: 245 WTVPSSA---CD--PSVHGALHITLSFTYMNSMLDPLVYFSSPSFPKFYNKLKICSLKP 299
          W + +S  C+  SV A  ITLSETYMNSMLDP+VYFSSPSFP F++ L  L+
Sbjct: 256 WLLHTSGTQNCVYRSVDLAFFITLSFTYMNSMLDPVVYFSSPSFPNFFSTLINRCLQR 315

Query: 300 KQPGHSKTQRPEEMPISNLGRRSCISVANSFQSQSDGQWDP 340 (residues 5-340 of SEQ ID NO:2)
          K G  R  + ++  ++  + + S W P
Sbjct: 316 KMTGEPDNNRSTSVELTGDPNKT-RGAPEALMANSGEPPWSP 355 (SEQ ID NO:4)

```

Hmmer search results:

Protein Family	amino acid residues
7 transmembrane receptor (rhodopsin)	32-278

4/5

FIGURE 3, page 1 of 2

```
1  AAATGACCAC  TTTTGCAAAA  TTGCATGCAT  TTCCAAGCTT  CATCCGGCTC
51  CAGGCTTGGC  CTCTCCCAGA  GGCAGGCGGC  TTGTGAGACG  GGCTCCAGAG
101 AAAGGACCTC  CCTGGGTCTC  TCATTTCCTG  GCTGAAGTTT  CTCTTCTCGC
151 TGCTGTGGCA  GCATCCAACC  CACACACACA  GGACCCGCAT  CCTGGGTGAT
201 GAAGTCAGAC  ACGCAGCAGC  TGGGTGAGTG  CTAACGCTCA  GATAAGCATC
251 TGTGCCATTG  TGGGGACTCC  CTGGGCTGCT  CTGCACCCGG  ACACTTGCTC
301 TGTCCCCGCC  ATGTACAACG  GGTCTGTCTG  CCGCATCGAG  GGGGACACCA
351 TCTCCCAGGT  GATGCCGCCG  CTGCTCATTG  TGGCCTTTGT  GCTGGGCGCA
401 CTAGGCAATG  GGGTCGCCCT  GTGTGGTTTC  TGCTTCCACA  TGAAGACCTG
451 GAAGCCCAGC  ACTGTTTACC  TTTTCAATTT  GGCCGTGGCT  GATTTCCTCC
501 TTATGATCTG  CCTGCCTTTT  CGGACAGACT  ATTACCTCAG  ACGTAGACAC
551 TGGGCTTTTG  GGGGACATTCC  CTGCCGAGTG  GGGCTCTTCA  CGTTGGCCAT
601 GAACAGGGCC  GGGAGCATCG  TGTTCCTTAC  GGTGGTGGCT  GCGGACAGGT
651 ATTTCAAAGT  GGTCCACCCC  CACCACGCGG  TGAACACTAT  CTCCACCCGG
701 GTGGCGGCTG  GCATCGTCTG  CACCCTGTGG  GCCCTGGTCA  TCCTGGGAAC
751 AGTGTATCTT  TTGCTGGAGA  ACCATCTCTG  CGTGCAAGAG  ACGGCCGTCT
801 CCTGTGAGAG  CTTTCATCATG  GAGTCGGCCA  ATGGCTGGCA  TGACATCATG
851 TTCCAGCTGG  AGTTCTTTAT  GCCCTCGGC  ATCATCTTAT  TTTGCTCCTT
901 CAAGATTGTT  TGGAGCCTGA  GGCGGAGGCA  GCAGCTGGCC  AGACAGGCTC
951 GGATGAAGAA  GGCGACCCGG  TTCATCATGG  TGGTGGCAAT  TGTGTTCATC
1001 ACATGCTACC  TGCCCAGCGT  GTCTGCTAGA  CTCTATTTCC  TCTGGACGGT
1051 GCCCTCGAGT  GCCTGCGATC  CCTCTGTCCA  TGGGGCCCTG  CACATAACCC
1101 TCAGCTTCAC  CTACATGAAC  AGCATGCTGG  ATCCCCTGGT  GTATTATTTT
1151 TCAAGCCCCT  CCTTTCCTAA  ATTCTACAAC  AAGCTCAAAA  TCTGCAGTCT
1201 GAAACCCAAG  CAGCCAGGAC  ACTCAAAAAC  ACAAAGGCCG  GAAGAGATGC
1251 CAATTTCGAA  CCTCGGTTCG  AGGAGTTGCA  TCAGTGTGGC  AAATAGTTTC
1301 CAAAGCCAGT  CTGATGGGCA  ATGGGATCCC  CACATTGTTG  AGTGGCACTG
1351 AACAAGCAGA  CCAACAACAC  TGAGGAAGAT  AGAGTGGTGA  CTTAGAATTA
1401 ACTCGTGCTA  AGGGGTTCGG  GGCTTTGAAA  ATGCCACCCC  CCTTCTTAT
1451 TGCAAGACGG  CTTCTCGCAC  ATGAACTGCA  TCCTTCTCAT  TCTGTCGGAA
1501 ATGAAATTCA  CACAACATA  CCTTTTGGG  AGGTTCCAGT  TGATTGAAGT
1551 GAGTTGGCTG  CATTTTCTTA  TCTGATCACA  ATGGCAGGGG  ACAGAATGTG
1601 CATGGAGTGG  AGCATGTGTG  TGTGGGAGG  GGGGCTAGGA  ACTGCACAGC
1651 CCTTGTGTAA  TTTTCGTTGT  TTGTTTTTGT  TTTGAGACAG  AGTCTCACTC
1701 TGTGTCCCAG  GCTGGAGTGC  AGTGGCACAG  TCTCGGCTCA  CTGCAACCTC
1751 TGCCTCCCGG  GTTCAAGCAA  TTCTCCTGCC  TCAGCCTCCC  GAGTAGCTGG
1801 GATTAGAGGC  GCCAGCCAAC  ACACCCGGCT  AATTTTGTGA  TTTTGTAGT
1851 AGACAGGGTT  TTGCCATGTT  GGCCAGGCTG  GTCTCGAGCT  CCTGACCTCA
1901 GGTGATCCGC  CTGCCTTGCC  CTCCCAAAGT  GGTGGGATCA  CAGGCGTGAG
1951 CCACCGTGCC  CGGCCTCCCC  TGTGTCATTT  TAAATGGCTA  AGTAAATGGG
2001 TATATGTGTT  TGAATGGGCG  ATGTTCACTC  TCTTAGGGGC  TATGGGGCAG
2051 TTAGCAGCAT  TTCTATCCT  CTGACCTTAA  ATCATTCCTT  ATCTCAGAAA
2101 ACAGAAACCG  GGCTCAGTCA  ATCAATGCTT  TATTTCAGGC  CGAATGAGGC
2151 TCTTTAGATT  GGGATCTATT  GATCTATCAA  TTTTCATCTT  TACATTTCTT
2201 TGTACATCTG  TACATTTTGT  CCAAATGTAC  ATCTGTACGT  CTGTCATCAT
2251 TGTGACTTCC  TGGTAGCCCA  AGAAGAACAA  CAACAAAACA  ATCTGCTCTG
2301 ACCTTCTTCA  AATCTTTGTA  TTTCAAAGAA  G (SEQ ID NO:3)
```

FEATURES:

Start: 311

Stop: 1349

5/5

FIGURE 3, page 2 of 2

MAP POSITION FOR ALLELIC VARIATION:

#	SHGCNAME	CHROM#	LOD_SCORE	DIST.(cRs)
1	SHGC-64103	12	6.12	45
2	SHGC-56772	12	6.12	45
3	SHGC-34758	12	6.06	45